

# Approaches to Aquatic Invasive Species Integrated Pest Management Control and Common ID


Chris Page  
South Carolina  
Department of Natural Resources



Aquatic Nuisance Species  
Program

# Important First Step:

- Identify your problem plant
- Know your Goals for the Pond
- Know about your downstream use





# Know the Law:

- Know which options may be viable and cost efficient
- Use only herbicides approved by the U.S. EPA
- Apply at the proper, legal, rates and selectively to the target plant(s)
- Apply as carefully as possible.
- You may only treat your private pond
- To treat someone else's pond you must be a licensed Category 5 applicator

# Pond Management– What's your goal?

- Recreational fishing
- Swimming & boating
- Wildlife habitat / aesthetics
- Livestock water supply
- Aquaculture
- Storm water management





# Basic Types of Control

- Physical – mechanical devices; cookie cutters, harvesters, rakes, etc; lake level manipulation; drawdowns.
- Biological – living organisms; pathogens, insects, fish.
- Chemical – specially designed and approved herbicide products.
- Integrated – one or more of the above



# Integrated Control



Aquatic Nuisance Species  
Program



# Aquatic Plant Control – Physical

- Benthic barriers
- Drawdown
- Mechanical removal
- Dredging
- Dyes



# Mechanical Harvesters





# Mechanical Harvesters

- Advantages

- Site selective control
- Removes plants from water
- Immediate relief of problem plants
- Perceived as environmentally safe

- Disadvantages

- Very expensive (\$600-1,000/ac)
- Provides only short-term control
- Spreads plant fragments
- Captures fish and other aquatic organisms
- Extremely slow (1-2 ac./day), so limited control
- Requires disposal sites



# Lake Level Drawdowns



Aquatic Nuisance Species  
Program



# Drawdowns

- Advantage

- Proven effective on some submersed vegetation
- Provides moderate to long-term control
- Controls plant growth over a large area
- Very inexpensive (little or no cost)

- Disadvantages

- Limits recreational uses (boating and swimming)
- Impairs homeowner access to lake or pond
- May require removal of larger boats from pond
- Requires advanced planning



# Sterile Grass Carp - *Ctenopharyngodon idella*



# Sterile Grass Carp

- Advantages
  - Long-term control
  - Relatively inexpensive (\$60-90/ac initial; \$12-16/ac after 5 years)
  - No water use restrictions
  - Can control plant growth over large areas
  - Do not reproduce
- Disadvantages
  - Slow control response
  - Impact non-target plant species
  - Difficult to regulate amount of control
  - Cannot control feeding location
  - Difficult to contain in water body





# Alligatorweed Flea Beetle - *Agasicles hygrophila*



# Alligatorweed Flea Beetle

- Advantages

- Long-term control
- Relatively inexpensive
- No Impact to non-target plant species
- No water use restrictions
- Can control plant growth over large areas

- Disadvantages

- Slow control response
- Difficult to regulate amount of control
- Cannot control feeding location
- Can be killed off by colder temperatures





# Aquatic Plant Control – Chemical



Aquatic Nuisance Species  
Program



# Aquatic Plant Control – Chemical

- Identify the problem plant
- Use only EPA registered and approved products
- Read and follow all label directions
- Timing
  - late spring, early summer
- Temperature
  - Over 65°
- Retreat?



# Aquatic Herbicides

- Advantages

- Effective on a range of plants
- Provides site selective control
- Limited water use restrictions
- Most are approved for drinking water supplies
- Can control relatively large areas (~80 ac/day)

- Disadvantages

- Controls some non-target species
- May be toxic to fish
- Some are slow acting
- Control level variable
- Limited effectiveness in deep water (>6 ft)



# Recent Advancements in Herbicides





# Recent Advancements in Herbicides

- Hardball, Sinkerball - Liquid 2, 4-d
  - \$30-40 per gallon - Up to 5 gallons per application (\$200)
- Vs granular 2, 4-d
  - \$2.50-3.50 per pound - 150-250 lbs per acre per application(\$875)
- Renovate Max-G - Mix of Renovate 3 and 2, 4-d
  - \$2.80-3.50 per pound - 150-200 lbs per acre per application(\$700)
  - Gives longer lasting control



- Renovate 3 - triclopyr
- Triclopyr, rapidly enters through the target plant's leaves and stems, interfering with plant metabolism, and providing systemic control of susceptible plant species. The herbicidal power of Renovate impacts most dicot (broadleaf) plants, while having little to no impact on most monocots (grassy type species), providing an excellent tool for aquatic ecosystem restoration programs.
- Renovate carries no restrictions on recreational use such as swimming and fishing, or on livestock consumption of water from the treatment area. Renovate can be used near active potable water intakes.



- **Galleon SC** - Penoxsulum
- Key target aquatic weed species: hydrilla, water hyacinth, water lettuce, salvinia species, frog's bit and duckweed.
- Multiple forms of application give flexibility to choose best management option:
- Injection for large-scale, in-water treatment of submersed and floating species
- Foliar sprays via ground, boat, or aerial application for targeted treatments of floating and emergent species
- Direct application to exposed littoral sediment for pre-emergence control following drawdown or other low-water events.





- **Clearcast - Imazamox**
- Cost - \$175 per gallon
- **Recommended Rates – Cattails**
- 64 ounces/acre broadcast
- Use rates 32-64 ounces/acre broadcast
- 1% solution spot spray
- **Recommended Rates – Chinese tallow**
- 64 ounces/acre broadcast
- 2% v/v solution spot spray
- 50% v/v solution hack & squirt
- 50% v/v solution cut stump



- **Clearcast Benefits**
- Applications can be made near desirable hardwood vegetation
- Controls the entire plant, including root system
- Minimal soil residual allows for re-colonization of desirable species
- Minimal irrigation restrictions
- Labeled for golf course waters used for irrigation



# Common Problems



Aquatic Nuisance Species  
Program



# Floating, not rooted



Aquatic Nuisance Species  
Program



# Water hyacinth - *Eichhornia crassipes*

- Thick, waxy, glossy leaves
- Stalks are bulbous and spongy
- Distinctive White and Bluish “orchid like” flower with 6 petals







Aquatic Nuisance Species  
Program



# Giant salvinia - *Salvinia molesta*



The upper surface is covered with dense, stiff white hairs with distinct “egg beater” shaped tips.








Aquatic Nuisance Species  
Program



# Water lettuce - *Pistia stratiotes*

- Thick, soft light green leaves; usually with small soft hairs covering the plant
- “Velvety” to the touch
- Resembles a small head of lettuce



# Water chestnut - *Trapa natans*




Aquatic Nuisance Species  
Program



# Watermeal

## *Wolffia columbiana*



Water meal  
*Wolffia columbiana*  
Photo by Vic Ramey  
Copyright 2000 Univ. Florida



Water meal  
*Wolffia columbiana*  
Photo by A. Murray  
Copyright 2000 Univ. Florida



Aquatic Nuisance Species  
Program



# Duckweed

## *Lemna minor*



Aquatic Nuisance Species  
Program





Aquatic Nuisance Species  
Program



# Common reed - *Phragmites australis*

- Alternate leaves
- Grows in clumps
- Bluish-green color
- Very dense growth habit
- 10-12 feet tall







Aquatic Nuisance Species  
Program





Aquatic Nuisance Species  
Program



# Water primrose - *Ludwigia hexapetala*

- Alternate willow-like leaves
- Bright yellow flower normally with 5 petals
- Can have different characteristics in rosette stage







Aquatic Nuisance Species  
Program



- Opposite willow-like leaves
- White “sno-cone” flower
- Branched, hollow stems







Aquatic Nuisance Species  
Program



# Alligatorweed

- Emergent, marginal plant
- Opposite entire leaves and hollow stems
- White, clover-like flower
- Spread by seed and fragmentation







# Alligatorweed

*Alternanthera philoxeroides*



Aquatic Nuisance Species  
Program





# Parrot feather

*Myriophyllum aquaticum*



Aquatic Nuisance Species  
Program



# Parrotfeather

- Submersed or emergent
- Feather-like leaves in whorls
- Grayish appearance to emergent foliage
- Stems may have reddish appearance
- Common throughout SC





# Submersed




Aquatic Nuisance Species  
Program

## Hydrilla - *Hydrilla verticillata*

- Rough to the touch
- Sharply toothed leaf margins, usually evident without magnification
- Usually 1-12 small, sharp teeth or spines on the lower midrib of the leaf
- Small, inconspicuous white pistillate flowers, less than 1/4 inch wide
- Number of leaves per whorl usually about the same at branching and non branching nodes; however, they may be double
- Leaves in whorls of 3-8







Aquatic Nuisance Species  
Program



# Brazilian elodea - *Egeria densa*



Aquatic Nuisance Species  
Program



# Slender naiad - *Najas minor*



Aquatic Nuisance Species  
Program





Aquatic Nuisance Species  
Program



*Utricularia biflora*



Photo by Jess Van Dyke  
Copyright 1998 Florida Department of Environmental Protection

# Submersed Bladderwort *Utricularia* spp.

*Utricularia floridana*



*Utricularia* spp.  
1996 Kerry Dressler



Aquatic Nuisance Species  
Program






Aquatic Nuisance Species  
Program





Aquatic Nuisance Species  
Program





Coontail  
*Ceratophyllum demersum*  
 Photo by V. Ramey  
 Copyright 2002 Univ. Florida

# Submersed Coontail *Ceratophyllum demersum*



*Ceratophyllum demersum*  
 1996 Kerry Dressler



Aquatic Nuisance Species  
 Program





Aquatic Nuisance Species  
Program





Two-leaf watermilfoil  
*Myriophyllum heterophyllum*  
 Photo by A. Murray  
 Copyright 2002 Univ. Florida

# Submersed Variable-leaf milfoil *Myriophyllum heterophyllum*



Two-leaf watermilfoil  
*Myriophyllum heterophyllum*  
 Photo by Vic Ramsey  
 Copyright 2002 Univ. Florida



Two-leaf watermilfoil  
*Myriophyllum heterophyllum*  
 Photo by A. Murray  
 Copyright 2002 Univ. Florida



Aquatic Nuisance Species  
 Program



# Submersed Bog moss *Mayaca fluviatilis*



*Mayaca fluviatilis*

Photo by Vic Ramey  
Copyright 1998 University of Florida



*Mayaca fluviatilis*

Photo by Vic Ramey  
Copyright 1998 University of Florida



Aquatic Nuisance Species  
Program



# Pondweeds



Waterthread pondweed  
*Potamogeton diversifolius*  
Photo by A. Murray  
Copyright 2001 Univ. Florida



Curly pondweed  
*Potamogeton crispus*  
Photo by Vic Ramey  
Copyright 2001 Univ. Florida

## *Potamogeton* species

**There are about 80 species of pondweeds in the world.  
Pondweeds are very important as wildlife food.**



Aquatic Nuisance Species  
Program





# Submersed Pondweeds

*Potamogeton sp.*



Aquatic Nuisance Species  
Program



# Variable-leaf pondweed, *Potamogeton diversifolius*



Aquatic Nuisance Species  
Program



*Potamogeton illinoensis*



Photo by Jess Van Dyke  
Copyright 1998 Florida Department of Environmental Protection

# Submersed Pondweeds

*Potamogeton illinoensis*

*Potamogeton illinoensis*  
1996 Alison Fox



*Potamogeton illinoensis*



Aquatic Nuisance Species

Program

Photo by Jess Van Dyke  
Copyright 1998 Florida Department of Environmental Protection




*Najas guadalupensis*



Photo by Brian Nelson  
Copyright 1998 Florida Department of Environmental Protection

## Submersed Southern Naiad *Najas guadalupensis*

*Najas guadalupensis*  
1996 Kerry Dressler



*Najas minor*




Photo by Jess Van Dyke  
Copyright 1998 Florida Department of Environmental Protection



Aquatic Nuisance Species  
Program



*Eleocharis* spp.



# Submersed to Emergent Spikerush *Eleocharis baldwinii*



*Eleocharis*



*Eleocharis baldwinii*



Aquatic Nuisance Species  
Road Grass  
Program

Photo by Ann Murray  
Copyright 1999 University of Florida

Photo by Ann Murray  
Copyright 1998 University of Florida



# Proliferating spikerush

## *Eleocharis baldwinii*



Aquatic Nuisance Species  
Program



# Floating Leaf

**Plants rooted in bottom,  
Most leaves float on the surface,  
or may be slightly raised above the surface  
in mature plants.**





# Watershield


## *Brasenia schreberi*



Water shield  
*Brasenia schreberi*  
Photo by Vic Ramey  
Copyright 1999 Univ. Florida



Water shield  
*Brasenia schreberi*  
Photo by Vic Ramey  
Copyright 1999 Univ. Florida



Water shield  
*Brasenia schreberi*  
Photo by Ann Murray  
Copyright 1999 Univ. Florida





# Fragrant water lily

## *Nymphaea odorata*



Fragrant water lily  
*Nymphaea odorata*  
Photo by A. Murray  
Copyright 2000 Univ. Florida



Aquatic Nuisance Species  
Program



# American lotus

## *Nelumbo lutea*



American lotus  
*Nelumbo lutea*  
Photo by V. Swaney  
Copyright 2001 Univ. Florida



American lotus  
*Nelumbo lutea*  
Photo by A. Murray  
Copyright 2001 Univ. Florida





# Spatterdock, cow lily

## *Nuphar lutea*



Spatterdock, cow lily  
*Nuphar advena*  
Photo by Vic Ramey  
Copyright 2000 Univ. Florida



*Nuphar advena*  
Photo by Vic Ramey  
Copyright 2005 Univ. Florida



Aquatic Nuisance Species  
Program



# Big floatingheart Banana lily

*Nymphoides aquatica*

Don't confuse with *Nymphaea mexicana* (yellow water lily)



Banana lily  
*Nymphoides aquatica*  
Photo by Vic Ramey  
Copyright 2001 Univ. Florida



*Nymphoides aquatica*  
tubers



Banana lily  
*Nymphoides aquatica*  
Photo by A. Murray  
Copyright 2001 Univ. Florida



Aquatic Nuisance Species  
Program



# Yellow floating heart

## *Nymphoides peltata*



Yellow floating heart  
*Nymphoides peltata*  
Lake Champlain Sea Grant/VIDEC  
photo by M. Malchoff



Yellow floating heart  
*Nymphoides peltata*  
Photo by Vic Ramey  
Copyright 2002 Univ. Florida



Aquatic Nuisance Species  
Program



# Web sites for more information

**SCDNR – Aquatic Nuisance Species Program**

**<http://www.dnr.sc.gov/invasiveweeds/>**

**SCDNR – Freshwater Fisheries**

**<http://www.dnr.sc.gov/water/aquaff/>**

**UF/IFAS Center for Aquatic and Invasive Plants**

**<http://plants.ifas.ufl.edu/>**



Aquatic Nuisance Species  
Program